

Simpson's Theory and Superrigidity of Complex Hyperbolic Lattices

ALEXANDER REZNIKOV

Abstract We attack a conjecture of J. Rogawski: any cocompact lattice in $SU(2, 1)$ for which the ball quotient $X = B^2/\Gamma$ satisfies $b_1(X) = 0$ and $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$ is arithmetic. We prove the Archimedean superrigidity for representation of Γ is $SL(3, \mathbb{C})$.

Théorie de Simpson et superrigidité des réseaux hyperboliques complexes

Résumé Soit $\Gamma \subset SU(2, 1)$ un réseau cocompact et soit $X = B^2/\Gamma$. Nous prouvons: si $b_1(X) = 0$ et $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$ alors tous les représentations ρ de Γ dans $SL(3, \mathbb{C})$ sont conjugué à la représentation naturelle ou la fermeture de Zariski de l'image $p(\Gamma)$ est compacte.

Version française abrégée - Le théorème classique de Margulis dit que tous les réseaux dans les groupes de Lie semi-simples sont superrigides. Ceci a été généralisé par Corlette [C] à la superrigidité des réseaux quaternioniques et de Cayley. D'autre part, Johnson et Millson ont montré qu'il existait des déformations des réseaux cocompact dans $SO(n, 1)$ si on regarde $SO(n, 1)$ comme plongé dans $SO(n + 1, 1)$.

C'est une question d'intérêt fondamental de savoir si les réseaux hyperboliques complexes sont superrigides.

Dans cet article, nous considérons la question suivante de J. Rogawski.

Hypothèse Soit $X = B^2/\Gamma, \Gamma \subset SU(2, 1)$ une surface hyperbolique complexe compacte. Supposons $b_1(X) = 0$ et $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) = \mathbb{Q}$. Alors Γ est arithmétique et provient d'une algèbre avec division $E|\mathbb{Q}$ de rang 3 avec une involution.

Observons que pour tous les réseaux provenant d'algèbres avec division, on a effectivement $b_1(X) = 0$ et $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) = \mathbb{Q}$ [Rog].

Soit ℓ un fibré linéaire tautologique sur X [Re]. La condition $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) = \mathbb{Q}$ dit que $[\ell] = k \cdot$ générateur dans $Pic(X)/tors \approx \mathbb{Z}$.

Le résultat principal de cet article prouve la superrigidité des représentations de Γ dans $SL(3, \mathbb{C})$ dans le cas $k = 1$.

Theoremé principal Soit $X = B^2/\Gamma$ et supposons que $b_1(X) = 0$ et $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) = \mathbb{Q}$. Soit $[\ell]$ un générateur de $Pic(X)/tors \approx \mathbb{Z}$. Si ρ est une représentation de Γ dans $SL(3, \mathbb{C})$ alors soit ρ est conjugué à la représentation naturelle de Γ , soit la fermeture de Zariski de l'image $\rho(\Gamma)$ est compacte.

Je vaudrais remercier Ron Livné, Jon Rogawski, et Carlos Simpson pour beaucoup de discussions intéressantes. Je vaudrais aussi remercier Marina Ville et Lucy Katz pour son aide essentielle à la préparation de cet article.

Simpson's Theory and Superrigidity of Complex Hyperbolic Lattices

ALEXANDER REZNIKOV

0 Main Theorem The classical theorem of Margulis establishes the superrigidity of lattices in semisimple Lie groups of rank ≥ 2 . The work of Corlette [C] extended this to (Archimedean) superrigidity of uniform quaternionic and Cayley lattices. On the other hand, by Johnson and Millson [JM] some uniform lattices in $SO(n, 1)$ admit deformations as mapped to $SO(n + 1, 1)$.

It is therefore of fundamental interest to study to what extent the complex hyperbolic lattices are superrigid. Since there are nontrivial holomorphic maps between different ball quotients [DM] one should confine oneself's look to lattices (or manifolds) "minimal" in some sense.

The present note addresses the following conjecture of Jon Rogawski.

Conjecture. Let $X = B^2/\Gamma, \Gamma \subset SU(2, 1)$ be a compact ball quotient. Suppose $b_1(X) = 0$ and $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$. Then Γ is arithmetic and comes from a division algebra $E|\mathbb{Q}$ of rank 3 with an involution.

Observe that for all lattices coming from division algebras, indeed $b_1(X) = 0$ and $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$ [Rog].

Let ℓ be the tautological line bundle over X [Re]. Since $Pic(X)/tors \approx \mathbb{Z}$, we have $[\ell] = k \cdot$ generator for some $k \in \mathbb{Z}$.

The main result of the paper establishes the superrigidity of representations of Γ in $SL(3, \mathbb{C})$ for Γ yielding $k = 1$ as follows.

Main Theorem. Let $X = B^2/\Gamma$ and suppose $b_1(X) = 0$ and $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$. If $[\ell]$ generates $Pic(X)/tors \approx \mathbb{Z}$, then any representation of $\Gamma = \pi_1(X)$ in $SL(3, \mathbb{C})$ is either conjugate to the natural representation up to the twist by a character, or has a compact Zariski closure.

One hopes, that, applying methods of [GS] one is able to prove the p -adic superrigidity and to settle Rogawski's conjecture.

I wish to thank Ron Livne, Jon Rogawski and Carlos Simpson for stimulating discussions.

1. Computations of Higgs bundles. We admit a knowledge of Simpson's theory [S1]. Let X be as above and let $\rho_0 : \Gamma \rightarrow PSU(n, 1)$ be the natural representation. Then the corresponding Higgs bundle is as follows [Re]. Take $E = TX \otimes \ell \oplus \ell$ as a holomorphic bundle and define $\theta \in H^0(\Omega^1 \otimes End(E))$ by $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$.

In view of the Simpson's theory, for proving the Main Theorem one needs to show that any complex variation of Hodge structure [S1] of type (2,1) over X is as above. Indeed, any representation is deformable to one, corresponding to a variation of Hodge structure [S2], and the natural representation is rigid [W].

So let $F = (\xi \oplus \eta, \theta)$ be a variation of complex Hodge structure, rank $\xi = 2$, rank $\eta = 1$, $\theta \in H^0(\Omega^1(X) \oplus Hom(\eta, \xi)) \approx H^0(Hom(TX \otimes \xi, \eta))$.

1.2. LEMMA. *Let λ, μ be rank two bundles over X and let $f \in H^0(Hom(\lambda, \mu)), f \neq 0$. Then either rank $f \leq 1$ everywhere or*

$$(c_1(\mu), [\omega]) \geq (c_1(\lambda), [\omega])$$

with the equality iff $\lambda \approx \mu$, and rank $f = 2$ everywhere. Here $[\omega]$ is the Kähler class.

PROOF: Consider $\wedge^2 f : \wedge^2 \lambda \rightarrow \wedge^2 \mu$. If $\wedge^2 f \neq 0$, then $\wedge^2 \mu \oplus (\wedge^2 \lambda)^{-1}$ has a nontrivial holomorphic section, whose zero locus is an effective divisor, so $(c_1(\wedge^2 \mu \otimes (\wedge^2 \lambda)^{-1}), [\omega]) \geq 0$ and the equality implies $\wedge^2 f$ is an isomorphism.

2. PROOF OF THE MAIN THEOREM:

Let $F = (\xi \oplus \eta, \theta)$ be as above.

Case 1 Rank $\theta = 2$ somewhere.

Applying the lemma, we get

$$(c_1(TX \otimes \xi), [\omega]) \leq (c_1(\eta), [\omega])$$

Now, $[\omega] \sim [\ell]$ since X is hyperbolic, and $c_1(TX) = -3[\ell]$ in $H^2(X, \mathbb{R})$, so

$$(c_1(\eta) - 2c_1(\xi), [\ell]) \leq 3[\ell]^2.$$

On the other hand, since $\xi \oplus \eta$ is a deformation of the flat bundle, $c_1(\xi \oplus \eta) = 0$, i.e. $c_1(\xi) = -c_1(\eta)$, so

$$(*) \quad (c_1(\xi), [\ell]) \leq [\ell]^2.$$

Since F is θ -stable [S1], $(c_1(\eta), [\ell]) < 0$, so $(c_1(\eta), [\ell]) > 0$. This leaves the only possibility $(c_1(\xi), [\ell]) = [\ell]^2$, because $[\ell]$ generates $Pic(X)/tors$. So $\xi = \ell \otimes \alpha$, where α is a linear unitary flat bundle, corresponding to $Pic(X)/tors \approx H_1^{tors}(X, \mathbb{Z})$. (recall that $b_1(X) = 0$). Moreover, since $(*)$ becomes an equality, we get by lemma above $\eta \approx TX \otimes \xi$ and θ takes the form $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$. Hence $F \approx (TX \otimes \ell \oplus \ell) \otimes \alpha$ and the proof is complete in this case.

Case 2 Rank $\theta \leq 1$ everywhere on X . There exists a collection of points (p_1, \dots, p_k) such that $\text{Ker } \theta$ extends to a rank one subbundle of $TX \otimes \xi$, say $\alpha \otimes \eta$. Since $H^2(X - \{p_1, \dots, p_k\}) \approx H^2(X)$, $c_1(\alpha)$ is well-defined in $H^2(X, \mathbb{Z})$. Moreover, by the removing of singularities in codimension two we have $H^i(X, \mathcal{O}) \approx H^i(X - \{p_1 \dots p_k\}, \mathcal{O})$, so from the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O} \rightarrow \mathcal{O}^* \rightarrow 1$ and the five-lemma we deduce that $H^1(X, \mathcal{O}^*) \approx H^1(X - \{p_1, \dots, p_k\}, \mathcal{O}^*)$, so $c_1(\alpha)$ is in the image of $Pic(X)$ in $H^2(X, \mathbb{Z})$. Let C be an irreducible curve of sufficiently high degree, which does not meet p_1, \dots, p_k . Since $TX \otimes \ell \oplus \ell$ remains θ -stable on C [S1] we get $(c_1(\alpha \otimes \ell)|_C, [C]) < 0$. In view of $H^{1,1}(X) \cap H^2(X, \mathbb{Q}) \approx \mathbb{Q}$ we can rewrite this as $(c_1(\alpha), [\ell]) < -[\ell]^2$. Since $[\ell]$ generates $Pic(X)/tors$, this actually means $(c_1(\alpha), [\ell]) \leq -2[\ell]^2$. Now, $c_1(TX) = -3[\ell]$, so $(c_1(TX/\alpha), [\ell]) \geq -[\ell]^2$. On C we have an isomorphism

$$\theta|_C : TX|_C \otimes \alpha \otimes \xi \rightarrow \text{Im } \theta \subset \eta|_C.$$

Hence $(c_1(\text{Im } \theta), [C]) = (c_1(TX/\alpha) + c_1(\xi), [C]) \geq (c_1(\xi) - [\ell], [C])$. Since, again, $[\ell]$ generates $Pic(X)/tors$ and $(c_1(\xi), [\ell]) > 0$ we get $(c_1(\xi) - [\ell], [C]) \geq 0$. This contradicts the θ -stability of $\xi \oplus \eta|_C$, because $\text{Im } \theta|_C$ is θ -invariant. The proof is complete.

References

- [C] K. Corlette, *Archimedean superrigidity and hyperbolic geometry*, Annals of Math., **135**, (1991), 165–182.
- [DM] P. Deligne, G. Mostow, *Commensurabilities Among Lattices in $PSU(1, n)$* , Annals of Mathematical Studies, **132**, Princeton Univ. Press, 1993.

- [GS] M. Gromov, R. Schoen, *Harmonic maps into singular spaces and p -adic superrigidity for lattices in groups of rank one*, Publ.Math.IHES, **76** (1992), 165–246.
- [JM] D. Johnson and J. Millson, *Deformation spaces associated to compact hyperbolic manifolds*, in: Discrete Groups in Geometry and Analysis (papers in honor of G.D. Mostow), Progress in Math. **67**, (1987), 48–106.
- [Re] A. Reznikov, *All regulators of flat bundles are torsion*, Annals of Math., (1995), to appear.
- [Rog] J. Rogawski, *Automorphic Representation of Unitary Groups in Three Variables*, Sect. 15.3, Princeton Univ. Press, 1990.
- [S1] C. Simpson, *Higgs bundles and local systems*, Publ. Math. IHES 75 (1992), 5–95.
- [S2] C. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety I*, Publ.Math.IHES **79** (1994), 47–129.
- [W] A. Weil, *Discrete subgroups of Lie groups I, II*, Annals of Math., **72** (1960), 369–384, **75** (1962), 578–602.

Institute of Mathematics

Hebrew University

Givat Ram 91904

ISRAEL

email: simplex@sunset.huji.ac.il